



# Long baseline neutrino experiments

L. Chaussard

## ► To cite this version:

L. Chaussard. Long baseline neutrino experiments. Heavy Quarks and Leptons 2002 HQ&L 2002, May 2002, Vietri Sul Mare, Palerme, Italy. pp.427-435. in2p3-00012337

**HAL Id: in2p3-00012337**

**<https://hal.in2p3.fr/in2p3-00012337>**

Submitted on 26 Feb 2003

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



*6th International Workshop  
Heavy Quarks and Leptons  
Vietri, May 27<sup>th</sup> - June 1<sup>st</sup>, 2002*

## *Session 9 : $\nu$ oscillations*

### *Long BaseLine $\nu$ experiments*

*(  $\nu$ -wars , episode I to episode III )*


*Lionel Chaussard  
OPERA Collaboration  
Institut de Physique Nucléaire de Lyon*



## Prologue: Why do we need LBL $\nu$ experiments ?

- Two flavour oscillation:  $P(\nu_\mu \rightarrow \nu_\tau) \sim \sin^2(2\theta) \sin^2(1.27 \Delta m^2 L/E)$


control  $\nu$  flavour purity and fluxes  
choose distance and energy



$$\begin{aligned} * \sin^2(2\theta)_{\min} &\sim (L) / (M_{\text{det.}}^{1/2}) \\ * \Delta m^2_{\min} &\sim 1 / (L^{1/2} M_{\text{det.}}^{1/4}) \end{aligned} \quad \left| \rightarrow \text{vary } L, \text{ large } M_{\text{det.}} \right.$$


(ref.: A. Ereditato, P. Migliozi, La Rivista del Nuovo Cimento, vol.23 n.12 (2000) 1-136)

- Three flavour oscillation: 3 angles  $\theta_{ij}$   
2 mass differences  $\Delta m^2_{ij}$   
1 ~~CP~~ phase  $\delta$




$\nu$  and  $\bar{\nu}$

## Episode I: "Standard" neutrino beam experiments



## K2K: from KEK to SuperKamiokande

- "disappearance" experiment
- 12 GeV protons on Al target, 1.3 GeV  $\nu_\mu$  beam from  $\pi$  decay
- near detector @ 300 m: 1kT water Cerenkov  
+ SciFi tracker +  $\mu$  range detector
- far detector @ 250 km: 50 kT (22.5 kT fiducial vol.) water Cerenkov
- status: started March '99  
SK accident on 12<sup>th</sup> of November 2001




## K2K: from KEK to SuperKamiokande

- hep-ex/0110034 v1 - 17 Oct.2001: using  $\sim 0.4 \cdot 10^{20}$  p.o.t.

44 fully contained events observed


$64^{+6.1}_{-6.6}$  expected without oscillation

- no oscillation hypothesis disfavored at  $\sim 2\sigma$



## K2K: from KEK to SuperKamiokande


- expectations for  $10^{20}$  p.o.t. (3-4 years more data taking):





## NUMI/MINOS: from Fermilab to Soudan

- "disappearance" experiment
- 120 GeV protons on graphite or Beryllium target
- 1 to 25 GeV  $\nu_\mu$  beam (3 configurations)
- near detector @ 290 m: 0.98 kT
- far detector @ 730 km: 5.4 kT
- far detector (extracted from S. Childress at NBI 2002) :  
8m octagonal tracking calorimeter,  
486 layers of 2.54 cm Fe, 2 sections, each 15 m long,  
4.1 cm wide plastic scintillator strips, WLS fibers + P.M.,  
25 800 m<sup>2</sup> active detector planes,  
magnet coil  $\langle B \rangle \approx 1.3\text{T}$
- status: far detector operational by late 2003  
near detector complete and tested late 2004  
beam commissioning ~ end 2004/ early 2005









# NUMI/MINOS: from Fermilab to Soudan



- 3 energy configurations:







## NUMI/MINOS: from Fermilab to Soudan

- many plots available from: <http://www-numi.fnal.gov/blessed/index.html>
- including stat. and syst. errors , for the 3 energy configurations
- for CC energy test, NC/CC ratio,  $\nu_\mu \rightarrow \nu_e$ ,  $\nu_s$  ...



## CNGS: from CERN to GranSasso


- 400 GeV protons on graphite target
- 15 GeV  $\nu_\mu$  beam




- status: K. Elsener @ NBI 2002: "CNGS is on schedule but..."  
first beam in Spring 2006 ... ?

## CNGS1: the OPERA project

- "appearance" experiment
- far detector 1.8 kT @ 730 km
- target tracker: emulsions sheets + lead, plastic scintillators, WLS fibers + P.M.
- spectrometers: drift tubes + resistive plate chambers (RPC)
- status: approved by CERN and LNGS in February 2001, constr. starts end 2002, scheduled for 2006




## CNGS1: the OPERA project



- Components:
  - 2 supermodules
  - 31 walls / supermodule
  - 52 x 64 bricks / wall
  - 56 **emulsion films** / brick

- Principle of the detection:


scintillators allow to find the hit brick (~ 30 bricks / day), which is **scanned** by microscopes (~ 20 cm<sup>2</sup> / hour) to find the "**kink**" of the tau



## CNGS1: the OPERA project

- expected number N of tau events:  
(5 years run with 1.8 kT average target mass)  
 $\Delta m^2 = 1.3 \cdot 10^{-3} \text{ eV}^2 : N = 4$   
 $2.5 \cdot 10^{-3} \text{ eV}^2 : N = 10$   
 $4.0 \cdot 10^{-3} \text{ eV}^2 : N = 26$
- background:  $\sim 0.65$  event
- proba. of  $3\sigma$  significance for  $\Delta m^2 = 2.5 \cdot 10^{-3} \text{ eV}^2 : \sim 99\%$
- room for improvement  
(efficiencies, beam x1.5 (?) )  
(ref.: Plan for the construction of the OPERA detector, May 2002)

- sensitivity (exclusion limits)








## CNGS/ICARUS:

- Liquid Argon TPC, 300 T modules
- 3D reconstruction of tracks
- status: almost approved...

T600 half-module mounted  
in Pavia March 2000 -  
March 2001

full test run April-  
August 2001



installation and cosmic  
run at LNGS  
early 2003





## CNGS/ICARUS:

- real  $K \rightarrow \mu \rightarrow e$  from the test run  
(ref. J.Rico, hep-ex/0205028 13 May 2002, Moriond Proc.)









## CNGS/ICARUS:

- $\nu_\mu \rightarrow \nu_\tau$  "appearance": example of the electron channel



Cut at 2: 13  $\nu_\tau$  CC, 0.7  $\nu_e$  CC

(ref. LNGS-EXP 13/89 add.2/01, November 2001)




"event shape" likelihood based on visible energy, missing  $P_\tau$ ,  $P_\tau(e)$ ,  $P_\tau(h)$




## CNGS/ICARUS:

- sensitivities (ref. talk from A.Rubbia, NNN02, January 2002)


$\nu_\mu$  disappearance  
by CC energy test  
( $\Delta(L/E) \sim 30\%$ )




tau "appearance"  
by missing  $P_+$  studies  
3  $\sigma$  effect: 40 kT.year ??




## Episode II: "SuperBeams"





## SuperBeams:

- physics: essentially  $\nu_{\mu} \rightarrow \nu_e$   
push  $\theta_{\text{CHOOZ}}$  limit
- technology:
  - \* increase proton beam **intensity** for a high luminosity  $\nu$  beam
  - \* **off-axis** detectors to work at **low energy** ( $\sim \text{GeV}$ )  
and to reduce  $\nu_e$  contamination
- possibilities:
  - \* **Japan**: JHF-I ( 0.75 MW ) / JHF-II ( 4 MW )
  - \* **U.S.A.**: new 1 MW proton driver for the Fermilab Booster  
or x4 intensity of Fermilab Main Injector  
or Brookhaven AGS to Homestake or New Mexico
  - \* **Europe**: 4 MW proton linac at CERN ?



## JHF: from Tokai to SuperKamioKande

- JAERI: 50 GeV PS delivers  $3.3 \cdot 10^{14}$  p / 3.3s (0.75MW)  
to be upgraded to **4MW**
- Kamiokande: SuperK (22.5 kT) to be upgraded to **HyperK** (1000 kT)
- narrow band  $\nu_{\mu}$  beam  **$\sim 1$  GeV**
- status : first phase approved  
starts **2007**
- goals: measure...  
..."everything" ?

(ref.: hep-ex/0106019, 5 June 2001)



## JHF: from Tokai to SuperKamioKande

- $\nu_\mu$  disappearance:  
gain an order of magnitude


$$\delta(\Delta m_{23}^2) \sim 10^{-4} \text{ eV}^2$$

$$\delta(\sin^2(2\theta_{23})) \sim 0.01$$


- $\nu_\mu \rightarrow \nu_e$ : gain a factor 20

$$\sin^2(2\theta_{\mu e}) \sim 0.5 \sin^2(2\theta_{13}) > 0.003$$

- $\cancel{CP}$  phase  $\delta$  down to  $\sim 20^\circ$   
in the case of LMA solution






## Episode III: "Neutrino Factories"





## NuFact's:


- studies in US, Europe, Japan






## NuFact's or SuperBeams ?

- see for example hep-ph/0204352, 30 April 2002, P.Huber et al.



## NuFact's or SuperBeams ?

- in the same ref., discussion of CP violation measurements (assuming LMA)




## NuFact's or SuperBeams ?


- CP violation measurements

LMA  
area
   
 lower bound
   
 best fit
   
 upper bound

Sensitivity to CP-Violation at  $\delta_{CP} = +\pi/2$



## Epilogue (?):



2001 ...





2005 ?




2006 ?



2006 ?



2007 ?



2015 ?

